

AD-A184 372

DEVELOPMENT OF A DNA-BASED METHOD FOR DISTINGUISHING
THE MALARIA VECTORS. (U) EMORY UNIV ATLANTA GA DEPT OF
BIOLOGY V FINNERTY JUN 87 DAMD17-85-C-5104

1/1

UNCLASSIFIED

F/G 6/13

ML



END
9 87
0FIC



MICROCOPY RESOLUTION TEST CHART

AD-A184 372

DTIC EFILE COPY

(2)

AD -----

Development of a DNA-Based Method for Distinguishing the Malaria Vectors,
Anopheles gambiae from *Anopheles arabiensis*

Annual Report

Victoria Finney, Ph.D.

June 1987

Approved by:

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT CENTER
Ft. Detrick, Frederick, MD 21701-5010

Contract No. DAMD17-85-C-5184

For further information contact:
Dr. Victoria Finney

Medical Research Institute of Chemotherapy

1000 Rockville Pike, Bethesda, MD 20205

Contractor's Name: Medical Research Institute of Chemotherapy

DTIC
SELECTED
SEP 04 1987
S E D
RE

87 9 1 075

ADA 184372

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Emory University Biology Department		6b. OFFICE SYMBOL (If applicable)	
6c. ADDRESS (City, State, and ZIP Code) Atlanta, GA 30322		7a. NAME OF MONITORING ORGANIZATION	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION US Army Medical Research & Development Command		8b. OFFICE SYMBOL (If applicable)	
8c. ADDRESS (City, State, and ZIP Code) Ft. Detrick, Frederick, MD 21701-5102		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-85-C-5184	
11 TITLE (Include Security Classification) Development of a DNA-based method for distinguishing the malaria vectors <u>Anopheles gambiae</u> from <u>Anopheles arabiensis</u>		10 SOURCE OF FUNDING NUMBERS	
12 PERSONAL AUTHOR(S) Finnerty, Victoria, Ph.D.		PROGRAM ELEMENT NO 62770A	PROJECT NO 3MI- 62770A870
13a. TYPE OF REPORT Annual Report	13b. TIME COVERED FROM 86-6-15 TO 87-6-14	14 DATE OF REPORT (Year, Month, Day) 1987 June	15 PAGE COUNT .1
16 SUPPLEMENTARY NOTATION			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Recombinant DNA/Species Identification Anopheline Malaria Vectors	
FIELD 06	GROUP 13		
06	01		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) The Anopheles gambiae complex includes six morphologically identical species. Two of these (<u>A. gambiae</u> and <u>A. arabiensis</u>) are the primary African malaria vectors today. Since two or more of the species are commonly sympatric, epidemiological studies to determine the involvement of each in malaria transmission have been difficult. This report describes an attempt to develop a DNA probe to distinguish <u>A. gambiae</u> from <u>A. arabiensis</u> . The DNA probe is a segment of cDNA from <u>A. gambiae</u> which displays an RFLP when the two species are compared in their analysis. Thus far the probe has proven to be extremely sensitive and specific even with short exposures to fixative, simple agarose gel electrophoresis, and staining with ethidium bromide. <u>A. arabiensis</u> and <u>A. gambiae</u> are similar enough to distinguish by this method at room temperature for as long as one month at 4°C. We have also attempted to use the probe method on slides to identify a collection of adult female mosquitoes collected and used for species identification. This method was successful in identifying the species contained during DNA extraction.			
20 DISTRIBUTION AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> OTHERS		21 ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Mrs. Victoria Finnerty		22b. IF RPT AND ABSTRACT ARE DIFFERENT, EXPLAIN None	

AD -----

Development of a DNA-Based Method for Distinguishing the Malaria Vectors,
Anopheles gambiae from *Anopheles arabiensis*

Annual Report

Johns Hopkins University, B.L.

July 1987

Report No.

Contract No. DAMD17-85-C-5184
Task Order No. 1

Contract No. DAMD17-85-C-5184

SUMMARY

The *Anopheles gambiae* complex includes six morphologically identical species, two of which (*A. gambiae* and *A. arabiensis*) are the primary African malaria vectors today. Since two or more of the species are commonly sympatric, epidemiological studies to determine the involvement of each in malaria transmission have been difficult. This report describes our efforts to develop a DNA probe to distinguish *A. gambiae* from *A. arabiensis*. The DNA probe is a fragment of rDNA from *A. gambiae* which displays an RFLP when the two species are compared by Southern analysis. Thus far the probe has proven to be extremely sensitive since it can be used even with short exposures to diagnose single adult mosquitoes (or parts thereof) of both sexes. Larvae and pupae are similarly easy to distinguish. Specimens kept dessicated at room temperature for as long as nine months can be stored. We have demonstrated that the DNA probe method can also be readily used on dessicated abdomens, while the thoraces have been used for sporozoite analysis. Blood meal analysis is easily done from the protein pellet obtained during DNA extraction. The DNA probe method has been directly compared to the GDM isozyme method and no exceptions were found. The DNA probe method can diagnose a number of individuals bearing rare GDM alleles which cannot be scored enzymatically. Finally, the DNA probe method, when directly compared to the isozyme method, shows virtually complete agreement. The major limitation of this method is the need to label the probe. A unique restriction endonuclease fragment length polymorphism at nucleotide 1000 of the probe can be easily labeled with a standard kinase. This is followed by hybridization with ³²P-labeled riboprobe and results of the assay are determined by autoradiography. An advantage of this method is that no radioactive waste is generated which will reduce the cost of the procedure.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Avail and/or	
Not	Special
A-1	



FOREWORD

Studies with Recombinant DNA: The investigator has abided by the National Institutes of Health Guidelines for Research Involving Recombinant DNA Molecules (April 1982) and the Administrative Practices Supplements.

Emory University Biosafety File No. 142-85

TABLE OF CONTENTS

	Page
Front Cover	
DD Form 1473	
Title Page	
Summary	1
Foreword	2
Table of Contents	3
List of Tables and Figures	4
Body of Report	
Background	5
Rationale	5
Experiments and Results	
a) Isolation of diagnostic cDNA fragment	5
b) Stability of diagnostic fragment	5
c) General evaluation of diagnostic probe	6
d) Compatiblity of diagnostic probe	6
e) Temperature of probe and enzyme method	7
f) Comparison of probe and enzyme methods	7
References	12
List of Figures	13

LIST OF FIGURES AND TABLES

- Figure 1 Lambda Agr12 restriction map
- 2 Hybridization of pAgr12A to EcoRI digests of single dried female mosquitoes
- 3 Hybridization of pAgr12A to single dried male and female mosquitoes or mosquito abdomens.
- 4 Map of Kenya showing locations from which specimens were obtained
- 5 ODH electromorphs found in the Kenya field samples
- 6 Hybridization of the pAgr12A probe to EcoRI digests of field specimens

Table 1 Results of testing abdomens of *A. gambiae* complex mosquitoes collected in Asembo area of Kenya in October 1985

2 DNA probe and ODH isozyme analyses of *A. gambiae* complex mosquitoes from Kenya

3 Results of testing individual field specimens by DNA probe and cytogenetic methods.

ANNUAL REPORT

1. Statement of Problem Under Study: The two major malaria vectors, *A. gambiae* and *A. arabiensis* are morphologically indistinguishable (1,2). Yet biological studies indicate that these two sympatric species may not be equally involved in malaria transmission in those areas where they co-exist (3,4). Therefore, the resolution of a number of important epidemiological question concerning their role in malaria transmission is currently impossible. Epidemiological studies require a reliable means for species identification of individual field specimens. Moreover, these individuals must also be assayed for the presence of the malaria sporozoite. Presently, the only completely reliable means for species identification of adults is based upon examination of ovarian nurse cell polytene chromosomes (5). Alternative procedures based on enzyme electromorphs or those based on cuticular hydrocarbon profiles (5) are not reliable. Clearly, there are numerous reasons why neither enzyme variation nor HPLC are practical epidemiological tools for field specimens. Thus far, however, several reliable immunological procedures to assay sporozoites in dried field specimens have just been developed (7-11). Therefore, a very useful addition to these epidemiological tools would be a means of reliably identifying the species of individual dried mosquitoes. This report will discuss our current efforts which have resulted in the development of a reliable species assay.

2. Background: Many of the major malaria vectors are members of species complexes, for instance, *A. culicifacies* (12), *A. leucosphyrus* (13), and the *A. farauti* sibling series (14). In these complexes, as well as in the *A. gambiae* complex, reliable species identification of individuals is currently tedious and difficult. Since malaria continues to represent a major world health problem, epidemiological studies with these species is crucial.

Our studies focused on two sympatric species, *A. gambiae* and *A. arabiensis*. The proposal hypothesized that the genomic DNA of these two species currently differs in ways that would permit reliable species identification. In particular, we sought to develop a species differentiating assay based upon restriction fragment length polymorphism as detected by either heterologous or species-specific probes. During our first year of work we found that certain species specific sequences would be most useful for the assay we sought to develop, and therefore our efforts have focused upon these sequences rather than heterologous probes.

3. Rationale: A substantial body of evidence argues that $Ae.$ *gambiae* is different from *Ae.* *arabiensis* (see above). Given that we expect to find small differences between *Ae.* *gambiae* and *Ae.* *arabiensis*, then such differences would provide an excellent epidemiological tool. The main advantage of a PCR-based assay is its great sensitivity. If Southern analysis is to be used, temperatures dependent thermal profile should be examined, and PCR melting curves of *Ae.* *gambiae* and *Ae.* *arabiensis* will easily be obtained.

The first stage of the process is to identify the relevant documents. This can be done by scanning the entire document or by using a search function to find specific terms or phrases. Once the relevant documents have been identified, they can be reviewed and analyzed to determine the key information contained within them.

least 200 copies per genome arranged in a few large tandem assays (18). Therefore, the rDNA genes possess the ability to yield useful RFLPs as well as species-specific sequences, both of which would be the basis for a diagnostic assay.

4. Experiments and Results: Development of a single mosquito species assay used to distinguish *A. gambiae* from *A. arabiensis*. The strategy employed in this work was to quickly identify portion(s) of an rDNA gene in *A. gambiae* which were non-coding regions; i.e., the introns and spacers. Such non-coding DNA fragment(s) would then be the basis for further studies to determine whether they could reveal an RFLP in *A. arabiensis* DNA.

(i) Isolation of a diagnostic cloned rDNA fragment. An *A. gambiae* genomic library was screened with a *Sciara coprophila* rDNA clone (19) which contains one complete cistron. Thirty-two *A. gambiae* rDNA-containing phage were isolated and selected for further analysis. These clones were restricted with various enzymes and subjected to Southern analysis, in order to find nonconserved regions that might be used to reveal differences between the species. The blots were therefore probed with *Sciara* rDNA which is not expected to hybridize to fragments from the nonconserved regions. Restriction fragments from such regions (those not hybridizing to the *Sciara* probe) were then isolated from gels and used to probe genomic Southern blots of *A. gambiae* and *A. arabiensis* DNA. Clone XAge12, shown in Figure 1, was found to contain a 0.59kb EcoRI-SalI restriction fragment which consistently showed a different pattern of hybridization to *A. gambiae* versus *A. arabiensis* genomic DNA. The 0.59kb EcoRI-SalI fragment is very close to the 3' terminus of the 28S region of the mosquito rDNA cistron. Hybridization of the *Sciara* and *Calliphora* probes is very weak in this region, suggesting a low degree of conservation, yet this fragment is highly conserved among different geographic isolates of the three member species in the *A. gambiae* complex thus far examined. EcoRI-SalI genomic digests invariably show the 0.59kb fragment, and there is no evidence for detectable levels of inter-cistronic variation in either of these two restriction sites. In summary, the probe shows an unambiguous difference between members of the complex thus far examined.

the fragment of the diagnostic fragment to single dried reservoirs. The left fragment was sublimed into the Pilkington M5+ Plasmid Stripseller Drier in System 4 and this product, pfd-176, has been used to produce a large amount of a gentamicin complex chloride and dried molasses. A quantity enough to be used in the usual technique will come off the stripseller at one time. In the presence of a barium sulfate as a suspending agent (1%), the dried molasses solution is put into tubes and suspended with 10% NaOH and 10% NaClO (10%) and further dried by the Pilkington Drier. This is then washed with 10% NaClO (10%) and 10% NaOH (10%) and dried again. The final product is a white off-white granular complex salt containing 40% of the diagnostic fragment of λ and 10% molasses. The Pilkington Drier is a good way to dry the fragments because it is a very rapid process.




Fig. 1. λ -Agr12 Restriction map. The approximate locations of the 18S and 28S regions were determined by hybridization with heterologous *Sciara* rDNA (pBC2), kindly provided by S. Gerbi, and *Calliphora* rDNA (pKB-42 and pKB-33), kindly provided K. Beckingham. λ -Agr12 contains slightly more than 1 rDNA cistron, including the NTS. The dashed line indicates weak hybridization to the heterologous probes. The .59kb EcoRI-SalI restriction fragment which reveals a diagnostic restriction fragment length polymorphism between *A. gambiae* and *A. arabiensis* is shown as a darkened bar.




Fig. 2. Hybridization of pAgri2A to EcoRI digests of single dried female mosquitoes. Species and geographic origin of specimens are as follows: (1) *A. melas* (The Gambia), (2) *A. arabiensis* (Sudan, SENNAR colony), (3) *A. arabiensis* (Sudan, G/MAL colony), (4) *A. arabiensis* (Kenya), (5) *A. arabiensis* (Burkina Faso), (6) *A. gambiae* (Tanzania), (7) *A. gambiae* (Zanzibar), (8) *A. gambiae* (Kenya), (9) *A. gambiae* (Nigeria), (10) *A. gambiae* (The Gambia, G3 colony).

tion in *A. arabiensis* and *A. melas* are probably due to inter-cistronic variation in the spacer region.

In summary, specimens dessicated by a very simple method show no evidence of DNA degradation even when stored at room temperature for as long as nine months. Moreover, in other preliminary experiments we found that other life stages such as second instar larvae and pupae (and obviously both sexes) are readily scored by the DNA probe.

(iii). Genomic location of the diagnostic probe. Organization of the rDNA cistron appears to be the same in both males and females, as judged by Southern blots of male and female DNA. However, the intensity of hybridization of α Agr:2A to genomic Southern blots, as shown in figure 3, indicates that males have a smaller number of total copies, which is expected if the rDNA genes reside on the X chromosome. *A. gambiae-A. arabiensis* hybrid female mosquitoes reared in the laboratory contain both of the parental types of rDNA cistrons (Fig. 3). Male hybrids, on the other hand, show the cistron structure of the female parent, indicating that the rRNA genes are located primarily if not exclusively on the X chromosome. This finding directly associates the diagnostic probe with that part of the mosquito genome (the X chromosome) currently used as the basis for cytogenetic speciation.

(iv). Compatibility of the DNA probe method with the sporozoite assay, and blood meal analysis. In order to determine whether the probe could be used to assay single mosquitoes for the presence of the malaria parasite, we obtained a number of field specimens which had been dessicated for at least 14 months. The mosquitoes were cut so that Dr. Collins retained the head and thorax for the sporozoite assay (21) and we tested the abdomens. The results, shown in Table 1, indicate that the diagnostic probe can readily distinguish species of only part of a dried specimen. The proportions of gambiæ and anatensis in Ae. vexans which we found are similar to those found by other workers. Since these species are quite close and we did these experiments at a time when our DNA extraction procedure had not been optimized, there are an unacceptable number of undetectable individuals shown in Table 1. Since then, however, we have had few if any undetectable individuals from the specimens so treated.

A second important consideration for a diagnostic probe is whether it is compatible with blood meal analysis. The DNA extraction method described here allows blood meal analyses. A single mosquito is placed in thermal plastic, suspended in a 1.5 ml microtiter Eppendorf tube along with 30 µl of the peptide solution (18 mM histone, 1.5M EDTA, 1% SDS, 10% Tris, 50 µM EGTA). The mixture is incubated at 55°C for 1 h. After incubation, 400 µl of 8 M urea/0.5 M acetate buffer is added to extract DNA. After extraction, 400 µl of 100 mM EGTA/1% SDS/ethanol is added to the supernatant and centrifuged at 10,000 × g for 10 min. After discarding the supernatant, the DNA is precipitated overnight with 2.5 M ethanol and resuspended in 100 µl of 10 mM Tris, 1 mM EGTA. The precipitated supernatant is collected and 10 µl of 100 µM biotinylated DNA probe is added to the supernatant. Finally, the labeled DNA probe is added to the microtiter tube containing the DNA template and the tube is sealed with Parafilm®. The microtiter tube is then placed in a 60°C oven for 1 h.

and the other two were the same as the first. The last two were the same as the first, except that they contained 10% of the total amount of the first two.




Fig. 3. Hybridization of pAGr12A to single dried male and female mosquitoes or mosquito abdomens. Lane (1) *A. gambiae* female, (2) *A. gambiae* female (blood-fed), (3) *A. gambiae* female (abdomen only), (4) *A. gambiae* male, (5) *A. arabiensis* female, (6) *A. arabiensis* male, (7) *A. melas* female, (8) *A. melas* male, (9) *A. gambiae*-*A. arabiensis* hybrid female, (10) *A. gambiae*-*A. arabiensis* hybrid male, (11) *A. arabiensis*-*A. gambiae* hybrid female, (12) *A. arabiensis*-*A. gambiae* hybrid male. Female parent is listed first for all hybrids. DNA from a single abdomen is clearly more than sufficient to make a species identification. Furthermore, the presence of a blood meal in the abdomen does not significantly reduce DNA yield. Dessicated individual pupae and larvae (all instars except the first) can also be readily speciated.

Table 1. Result of testing abdomens of *A. gambia* complex mosquitoes collected in Asembo area of Kenya in October 1985.

Abdomens from:	Species		DNA not readable
	<i>A. gambia</i>	<i>A. arabiensis</i>	
Plasmodium falciparum infected mosquitoes	47 (75%)	17 (27%)	8
Uninfected mosquitoes	78 (49%)	80 (51%)	19

Note: percentages are based on specimens which were identified as to species. The sporozoite assay (21) and DNA probe assay were performed in December 1986.

lected by Dr. Collins in western Kenya (Ahero, Asembo, and Gumbi) and areas adjacent to the Lake Victoria area. Babaki, which are shown in figure 1, were used. These are relatively uninfested areas where the two species are known to be sympatric. The area is indicated, being a low-lying area on the high plain approximately 10 km east of the town of Kisumu. Previous studies of the mosquito fauna of this location have shown that most of the *A. gambiae* complex mosquitoes breeding in the area feed on humans (Bisanzio 1979). Asembo is a farming community located on the north shore of Lake Victoria, approximately 45 km west of Kisumu. Between Asembo and Gumbi is situated at a slightly higher elevation (approximately 200 m) the town of Kakamega. Gumbi terrain is considerably more hilly and mountainous than Babaki or the other areas in Ahero. Both *A. gambiae* and *A. strobli* are found throughout these three areas (Bisanzio 1979). Githuri is a small village located approximately 10 km west of Nairobi. During the summer months it has been referred to as Babati (Bisanzio 1979). Githuri is located in the same area as the other sites, but falls in the zone of a wetter environment (Bisanzio 1979). The weather in the area is made dry by a rainless period in March 1980, an intermediate climate state between the rainy season and the dry season. The following table summarizes the climatic data for each of the four areas. After completion of the field work, data was collected from the literature for each area, and the following information was obtained from the literature: rainfall, temperature, relative humidity, and wind speed.




Figure 4. Map of Kenya showing locations from which specimens were obtained.




Fig 5

Octanol dehydrogenase electromorphs found in the Kenya field samples. Lane 1, A. arabiensis from the G/MAL colony; lanes 2-3 are A. arabiensis from Ahero; lane 4, is an A. arabiensis (G/MAL) x A. gambiae (G3) hybrid produced in the laboratory; lanes 5-7, are A. gambiae from the GO-66 colony established with specimens collected in Gombe.




Figure 6 Hybridization of the pAg112A probe to EcoR1 digests field-collected specimens. Lanes 1-5, individual mosquitoes from different Ahero families; lanes 6-9, individuals from Gombe families.

Table 2. DNA probe and Odh isozyme analyses of *Anopheles gambiae* complex mosquitoes from Kenya.^a

Location	No.	Probe-Checked Families	ODH Alleles Present				
			Probe Result	90	95	98	100
Ahero	1	<u>A. gambiae</u>					+
	3	<u>A. arabiensis</u>		+	+		
	1	<u>A. arabiensis</u>			+		
	1	<u>A. arabiensis</u>		+	+		
Asembo	1	<u>A. gambiae</u>				+	+
	8	<u>A. gambiae</u>				+	
	1	<u>A. gambiae</u>			+	+	
	1	<u>A. arabiensis</u>		+	+	+	
	1	<u>A. arabiensis</u>			+		
	1	<u>A. arabiensis</u>		+	+	+	
	4	<u>A. arabiensis</u>		+	+		
	3	<u>A. arabiensis</u>		+			
Gombe	10	<u>A. gambiae</u>				+	
	1	<u>A. arabiensis</u>		+	+	+	
	1	<u>A. arabiensis</u>		+		+	
	1	<u>A. arabiensis</u>		+	+		+
	1	<u>A. arabiensis</u>		+	+		
	1	<u>A. arabiensis</u>		+	+		
Sabaki	86	<u>A. gambiae</u>				+	
	1	<u>A. gambiae</u>				+	+
	25	<u>A. arabiensis</u>		+			

^a Results for material from Ahero, Asembo, and Gombe represent analysis of at least two mosquitoes from each family for DNA type and an additional two mosquitoes for ODH isozymes. Results for material from Sabaki represent DNA probe and ODH isozyme analyses on single mosquitoes (the abdomen being used for ODH analysis and the head-thorax portion being used for DNA typing).

alleles test as *A. gambiae* by DNA probe; those with ODH-95 have the *A. arabiensis* DNA pattern.

Specimens from the western Kenya locations were treated somewhat differently in that individuals from each (sofemale) family were analyzed either by DNA probe (two individuals/family) or ODH isozyme (two or three individuals/family). Of the 41 different families so analyzed, none showed any within-family variation in the DNA probe hybridization pattern. Furthermore, only the expected 1.4Kt or 8Kb bands of hybridization were observed.

All the previous cited studies of ODH alleles in field specimens of *A. gambiae* and *A. arabiensis* indicate that, with near certainty, families with only the ODH-10 or ODH-15 alleles can be classified as *A. gambiae* and families with only the ODH-95 or ODH-96 alleles as *A. arabiensis*. Indeed, the 20 families from Kibera, Asembo, and Gomboni with the *A. gambiae* isozyme type show the DNA probe patterns diagnostic of *A. gambiae*. Also, the six families with only alleles ODH-95 or ODH-96 are identified by the probe as *A. arabiensis*. The fifteen families with other combinations of ODH alleles cannot be assigned to species on the basis of their isozyme pattern. However, the DNA probe test of these families indicates that 10 of the 15 are *A. arabiensis*, a finding that is consistent with those of Mulla, Thrusfield, and others who have reported considerably higher frequency of the isozyme 95 in collections of populations of *A. arabiensis* than *A. gambiae*. None of the remaining 5 individuals examined in this study gave a DNA probe reaction that would suggest an intermediate status.

The 54 of the 170 individual specimens from Kenya and the 36 families from Tanzania were analyzed where the probe results were not available; species diagnosis is therefore based on agreement with those predicted by the DNA probe. In addition, isozyme tests for ODH-95 were present in 36 individuals. In 30 of the 36, the ODH-95 pattern was determined, of these individuals, 27 had ODH-95 and 3 had ODH-96. The single individual with the ODH-95 pattern was from a family which had ODH-95 by DNA probe.

Thus, the 170 individual specimens from Kenya and the 36 families from Tanzania were analyzed where the probe results were available; species diagnosis is based on agreement with those predicted by the DNA probe. In addition, isozyme tests for ODH-95 were present in 36 individuals. In 30 of the 36, the ODH-95 pattern was determined, of these individuals, 27 had ODH-95 and 3 had ODH-96. The single individual with the ODH-95 pattern was from a family which had ODH-95 by DNA probe.

Table 3. Results of testing individual field specimens by DNA probe and cytogenetic methods.

COLLECTION SITE	CHROMOSOME RESULT	DNA RESULT		
		SAME AS CHROMOSOME	DIFFERENT FROM CHROMOSOME	NOT DONE OR NOT READABLE
ZIMBABWE	A. arab. (10)	10	0	0
	A. quad. (41)	41	0	0
KENYA				
Ahero	A. arab. (30)	76	0	4
	A. arab. (28)	25	1 (A. gamb.)	2
	A. gamb. (88)	70	1 (A. arab.)	17

References

- (1) Coluzzi M, A Sabatini, V Petrarca, and MA DiDeco. 1979. Chromosomal differentiation and adaptation to human environments in the Anopheles gambiae complex. *Trans R Soc Trop Med Hyg* 73:483-97.
- (2) Davidson G, HE Paterson, M Coluzzi, GF Mason, and DW Micks. 1967. The Anopheles gambiae complex. In: *Genetics of Insect Vectors of Disease* (ed. JW Wright and R Pal), Amsterdam, Elsevier, pp. 211-50.
- (3) Coluzzi M, A Sabatini, V Petrarca, MA DiDeco. 1977. Behavioral divergences between mosquitoes with different inversion karyotypes in polymorphic populations of the Anopheles gambiae complex. *Nature* 266:832-33.
- (4) Molibneaux L and G Gramiccia. 1980. The Garki Project: Research on the epidemiology and control of malaria in the Sudan savanna of West Africa. World Health Organization, Geneva.
- (5) Coluzzi M and A Sabatini. 1967. Cytogenetic observations on species A and B of the Anopheles gambiae complex. *Parasitologia* 9:73-88.
- (6) Carlson DA and MW Service. 1980. Identification of mosquitoes of Anopheles gambiae species complex A and B by analysis of cuticular components. *Science* 207:1089-91.
- (7) Zavala F, RW Gwadz, FH Collins, RS Nussenzweig, and V Nussenzweig. 1982. Monoclonal antibodies to circumsporozoite proteins identify the species of malaria parasite in infected mosquitoes. *Nature* 299:737-38.
- (8) Collins FH, F Zavala, PM Graves, AH Cochrane, RW Gwadz, J Akoh, and RS Nussenzweig. 1984. First field trial of an immunoradiometric assay for the detection of malaria sporozoites in mosquitoes. *Am J Trop Med Hyg* 33:538-43.
- (9) Collins FH, RW Gwadz, LC Koontz, F Zavala, and RS Nussenzweig. 1984. Laboratory assessment of a species-specific radioimmunoassay for the detection of malaria sporozoites in mosquitoes. *J Med Entomol* (in press).
- (10) Burkot TR, F Zavala, RW Gwadz, FH Collins, RS Nussenzweig, and DR Roberts. 1984. Identification of malaria-infected mosquitoes by a two-site enzyme-linked immunosorbent assay. *Am J Trop Med Hyg* 33:227-31.
- (11) Burkot TR, JL Williams, and I Schneider. 1984. Identification of Plasmodium falciparum infected mosquitoes by a direct antibody enzyme-linked immunosorbent assay. *Am J Trop Med Hyg* (in press).
- (12) Miles SJ. 1981. Unidirectional hybrid sterility from crosses between species A and species B of the taxon Anopheles (Cellia) culicifacies Giles. *J Trop Med Hyg* 84:13-16.
- (13) Baimai V, BA Harrison, and L Somchit. 1981. Karyotype differentiation of three anopheline taxa in the Balabacensis complex of Southeast Asia (Diptera: Culicidae). *Genetica* 57:81-86.
- (14) Mahon RJ and PM Miethke. 1982. Anopheles farauti No. 3, a hitherto unrecognized biological species of mosquito within the taxon A. farauti Laveran (Diptera: Culicidae). *Trans R Soc Trop Med Hyg* 76:8-12.
- (15) Jeffreys AJ. 1981. Recent studies of gene evolution using recombinant DNA. In: *Genetic Engineering 2* (ed R Williamson), NY, Academic Press, pp. 1-48.
- (16) Avise J, RA Lansman, and RO Shade. 1979. Use of restriction endonucleases to measure mtDNA sequence relatedness in natural populations. I. *Peromyscus*. *Genetics* 92:279-95.
- (17) Langley CH, E Montgomery, and W Quattlebaum. 1982. Restriction map variation in the AdH region of Drosophila. *Proc Nat Acad Sci USA* 79:5631-35.
- (18) Beckingham K. 1980. The ribosomal DNA of Calliphora erythrocephala: An analysis of hybrid plasmids containing ribosomal DNA. *J Mol Biol* 137:349-73.
- (19) Washino RK and CH Tempelis. 1983. Blood meal analysis. *Ann Rev Entomol* 28:179-202.

- (20) Livak K. 1984. Organization and mapping of a sequence on the *Drosophila melanogaster* X and Y chromosomes that is transcribed during spermatogenesis. *Genetics* 107:611-34.
- (21) Collins FH, F Zavala, PM Graves, AH Cochrane, RW Gwadz, J Akoh, and RS Nussenzweig. 1984. First field trial of an immunoradiometric assay for the detection of malaria sporozoites in mosquitoes. *Am J Trop Med Hyg* 33:227-30.
- (22) White GB. 1972. The Anopheles gambiae complex and malaria transmission around Kisumu, Kenya. *Trans R Soc Trop Med Hyg* 66:572-81.
- (23) Mosha FW and R Subra. 1982. Ecological studies on Anopheles gambiae complex sibling species in Kenya. I. Preliminary observations on their geographical distribution and chromosomal polymorphic inversions. WHO/VBC/82.867, 9pp.
- (24) Miles SJ. 1979. A biochemical key to adult members of the Anopheles gambiae group of species (Diptera: Culicidae). *J Med Entomol* 15:297-99.
- (25) Miles SJ. 1978. Enzymatic variation in the Anopheles gambiae Giles group of species (Diptera: Culicidae). *Bull Entomol Res* 68:85-96.

DISTRIBUTION LIST

12 copies

Director
Walter Reed Army Institute of Research
Walter Reed Army Medical Center
ATTN: SGRD-UWZ-C
Washington, DC 20307-5100

1 Copy

Commander
US Army Medical Research and Development Command
ATTN: SGRD-RMI-S
Fort Detrick, Frederick, Maryland 21701-5012

2 copies

Defense Technical Information Center (DTIC)
ATTN: DTIC-DDAC
Cameron Station
Alexandria, VA 22304-6145

1 copy

Dean
School of Medicine
Uniformed Services University of the
Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20814-4799

1 copy

Commandant
Academy of Health Sciences, US Army
ATTN: AHS-CDM
Fort Sam Houston, TX 78234-6100

END

q-87

DTIC